
























Drug names		Recommended adult dose *	Total daily pills
<b>Fixed dose combinations</b>			
0 5 10 15 20mm <i>Approximate to actual size.</i>			
Atripla (efavirenz + emtricitabine + tenofovir DF)		One tablet, once-daily. Take at night and not with a high fat meal. See info on separate drugs.	1
Biktarvy (bictegravir + TAF + emtricitabine)		One tablet, once-daily. Take with or without food. See info on separate drugs.	1
Eviplera (rilpivirine + emtricitabine + tenofovir DF)		One tablet, once-daily, with food (400 kcal). See separate drug info.	1
Odefsey (rilpivirine + emtricitabine + TAF)		One tablet, once-daily, take with food. See info on separate drugs.	1
Triumeq (dolutegravir + abacavir + lamivudine)		One tablet, once-daily. Take with or without food. See info on separate drugs.	1
Genvoya (elvitegravir + cobicistat + emtricitabine + TAF)		One tablet, once-daily. Take with food. See info on separate drugs.	1
Stribild (elvitegravir + cobicistat + emtricitabine + tenofovir DF)		One tablet, once-daily, take with food. See info on separate drugs.	1
Symtuza (darunavir + cobicistat + emtricitabine + TAF)		One tablet, once-daily, take with food. See info on separate drugs.	1
Juluca (dolutegravir + rilpivirine)		One tablet, once-daily, take with food. See info on separate drugs.	1

<b>Dual nukes: nucleoside or nucleotide reverse transcriptase inhibitors (NRTIs)</b>			
tenofovir DF 300 mg + emtricitabine 200 mg (Truvada [pictured] or generic) **		One tablet, once-daily.	1
TAF (10 mg white or 25 mg blue) + emtricitabine (200 mg), Descovy		One tablet, once-daily.	1
abacavir 600 mg + lamivudine 300 mg (Kivexa or generic) ***		One tablet, once-daily.	1
<b>Single nukes (NRTIs)</b>			
lamivudine (3TC) ** (Epivir [pictured] or generic)		1 x 300 mg (shown) or 2 x 150 mg, (taken as a once-daily or twice-daily dose).	1 if 300 mg 2 if 150 mg
abacavir ** (Ziagen [pictured] or generic)		2 x 300 mg tablets (taken as a once-daily or twice-daily dose).	2
emtricitabine (FTC) (Emtriva)		1 x 200 mg capsule, once-daily.	1
tenofovir DF * ** (Viread [pictured] or generic)		1 x 300 mg tablet, once-daily.	1

<b>NNRTIs: non-nucleoside reverse transcriptase inhibitors (non-nukes)</b>			
efavirenz 600 mg or 200 mg (Sustiva, [pictured] or generic) **		1 x 600 tablet (or 3 x 200 caps) once-daily; at night, not with high fat meal.	1 tablet (or 3 capsules)
nevirapine PR 400 mg (Viramune [pictured] or generic) **		1 x 400 mg once a day. Take with or without food.	1 x 400 mg
etravirine (100 mg or 200 mg) (Intelence)		2 x 100 mg OR 1 x 200 mg, twice daily, take with food. Dispersible in water.	2 or 4
rilpivirine (Edurant)		1 x 25 mg tablet, once-daily, take with main meal (500 kcal).	1

<b>INIs or INSTIs: integrase inhibitors</b>			
raltegravir 400 mg (pink) & 600 mg (yellow) (Isentress)		1 x 400 mg, twice-daily OR 2 x 600 mg tablet, once-daily. Take with or without food.	2
dolutegravir (Tivicay) *		1 x 50 mg tablet, once-daily (or 1 x 50 mg twice -daily). With food if twice-daily but with or without otherwise.	1 or 2
elvitegravir and bictegravir are only in combination pills – see Stribild, Genvoya and Biktarvy.			1

<b>CCR5 inhibitors (entry inhibitor)</b>			
maraviroc * (Celsentri)		150 mg or 300 mg or 600 mg, as directed, depending on other ARVs in the combination.	1 or 2 or 4

<b>b/PI: boosted protease inhibitors</b>			
atazanavir * (Reyataz)		1 x 300 mg cap + booster, once-daily. Take with food. 150 mg and 200 mg capsules also available.	1 (+ 1 booster)
darunavir 600 mg (orange) & 800 mg (red), (Prezista) *		1 x 800 mg + booster once-daily (OR 1 x 600 mg + 100 mg booster twice-daily with resistance). Take with food.	1 or 2 (+ 1 or 2 boosters)
atazanavir/cobicistat (Evotaz)		1 tablet, once-daily. Take with food.	1
darunavir/cobicistat (Rezolsta)		1 tablet, once-daily. Take with food.	1

<b>PK (pharmacokinetic) boosters</b>			
cobicistat (/c) (Tybost)		150 mg tablet, once daily. Used to boost atazanavir, darunavir and elvitegravir.	depends on boosted drug
ritonavir (/r) * (Norvir)		100 mg tablets used at different doses to boost other PIs.	depends on PI

\* Different doses or formulations might be used - always check doses with your pharmacist.

\*\* Generic versions might be a different colour and shape. § EU approval pending.